

Mid-Childhood Outcomes of Repeat Antenatal Corticosteroids: A Randomized Controlled Trial

Caroline A. Crowther, MD,^{a,b} Peter J. Anderson, PhD,^{c,d} Christopher J.D. McKinlay, PhD,^a Jane E. Harding, DPhil,^a Pat J. Ashwood, BSc(Hons),^b Ross R. Haslam, MBBS,^e Jeffery S. Robinson, MD,^b Lex W. Doyle, MD,^{c,d,f} for the ACTORDS Follow-up Group

abstract

OBJECTIVE: To assess if exposure to repeat dose(s) of antenatal corticosteroids has beneficial effects on neurodevelopment and general health in mid-childhood, at 6 to 8 years' corrected age.

METHODS: Women at risk for very preterm birth, who had received a course of corticosteroids ≥ 7 days previously, were randomized to intramuscular betamethasone (11.4 mg Celestone Chronodose) or saline placebo, repeated weekly if risk of very preterm birth remained. Mid-childhood assessments included neurocognitive function, behavior, growth, lung function, blood pressure, health-related quality of life, and health service utilization. The primary outcome was survival free of neurosensory disability.

RESULTS: Of the 1059 eligible long-term survivors, 963 (91%) were included in the primary outcome; 479 (91%) in the repeat corticosteroid group and 484 (91%) in the placebo group. The rate of survival free of neurosensory disability was similar in both groups (78.3% repeat versus 77.3% placebo; risk ratio 1.00, 95% confidence interval, 0.94–1.08). Neurodevelopment, including cognitive function, and behavior, body size, blood pressure, spirometry, and health-related quality of life were similar in both groups, as was the use of health services.

CONCLUSIONS: Treatment with repeat dose(s) of antenatal corticosteroids was associated with neither benefit nor harm in mid-childhood. Our finding of long-term safety supports the use of repeat dose(s) of antenatal corticosteroids, in view of the related neonatal benefits. For women at risk for preterm birth before 32 weeks' gestation, ≥ 7 days after an initial course of antenatal corticosteroids, clinicians could consider using a single injection of betamethasone, repeated weekly if risk remains.



^aLiggins Institute, University of Auckland, Auckland, New Zealand; ^bDiscipline of Obstetrics and Gynaecology, School of Medicine, The University of Adelaide, Adelaide, Australia; ^cClinical Sciences, Murdoch Children's Research Institute, Parkville, Victoria, Australia; ^dDepartment of Paediatrics, University of Melbourne, Parkville, Victoria, Australia; ^eThe Women's and Children's Hospital, Adelaide, South Australia, Australia; and ^fDepartment of Obstetrics and Gynaecology, The Royal Women's Hospital, University of Melbourne, Parkville, Victoria, Australia

Dr Crowther conceptualized and designed the study, obtained funding, coordinated and supervised data acquisition, carried out initial analyses and interpretation of data, and drafted the article; Dr Anderson conceptualized and designed the study, and participated in acquisition of data and interpretation; Dr McKinlay participated in coordination and acquisition of data, analyses, and interpretation; Dr Harding conceptualized and designed the study, obtained funding, coordinated and supervised data acquisition, and participated in interpretation; Mr Ashwood coordinated and supervised data acquisition; Dr Haslam coordinated data acquisition, and participated in data interpretation; Dr Robinson conceptualized and designed the study, and

WHAT'S KNOWN ON THIS SUBJECT: Repeat dose(s) of corticosteroids before very preterm birth reduces neonatal respiratory morbidity and serious outcomes in the newborn period, without evidence of harm at preschool age. The effect on health and neurodevelopment in later childhood is unknown.

WHAT THIS STUDY ADDS: Use of repeat dose(s) of antenatal betamethasone when indicated reduces neonatal morbidity without beneficial or adverse effects on survival free of neurosensory disability in mid-childhood, or evidence of any other effects.

To cite: Crowther CA, Anderson PJ, McKinlay CJ, et al. Mid-Childhood Outcomes of Repeat Antenatal Corticosteroids: A Randomized Controlled Trial. *Pediatrics*. 2016;138(4):e20160947

Repeat doses of corticosteroids before very preterm birth reduce neonatal respiratory morbidity and serious outcomes in the newborn period.¹ Systematic review of 4 trials²⁻⁵ that have reported on health outcomes in early childhood after repeat dose(s) of antenatal corticosteroids found no differences between exposed or unexposed children.¹ However, the efficacy and safety effects on health in later childhood are unknown.

In animals, repeat doses of antenatal corticosteroids have been associated with long-term adverse effects on blood pressure and brain mass,^{6,7} and a human observational study has reported behavioral problems.⁸

The Australasian Collaborative Trial of Repeat Doses of Corticosteroids for the Prevention of Neonatal Respiratory Disease (ACTORDS) reported clear evidence of neonatal benefit up to the time of hospital discharge, including reduced neonatal respiratory and serious morbidity.² Although *z*-scores at birth for weight and head circumference were reduced in infants randomly assigned to repeat corticosteroids compared with placebo, by hospital discharge, no significant differences were seen between groups.² Early childhood follow-up found no differences in survival free of major neurosensory disability between groups.⁹

Despite evidence of neonatal benefit and reassuring findings at early childhood follow-up, repeat corticosteroids are not commonly prescribed partly because of concerns about possible longer-term adverse effects on the child.

Therefore, we reassessed surviving children whose mothers participated in the ACTORDS at 6- to 8-years' corrected age. We hypothesized that exposure to repeat dose(s) of antenatal corticosteroids would have beneficial effects on the rate of survival free of any neurosensory disability, cognitive function,

behavior, educational achievement, body size, blood pressure, general health including respiratory health, and health-related quality of life.

METHODS

Design and Study Population

Eligible children were born to women recruited into the ACTORDS. As previously described,² women were eligible for the trial if at risk for preterm birth at <32 weeks' gestation with a singleton, twin, or triplet pregnancy, and they had received an initial course of antenatal corticosteroids ≥ 7 days prior.

Intervention

Women were randomized to either the repeat corticosteroid group (one 2-mL dose Celestone Chronodose, which contains 11.4 mg betamethasone, comprising 7.8 mg betamethasone sodium phosphate and 6 mg betamethasone acetate) or the saline placebo group. If the woman remained at risk for preterm birth, the allocated study treatment could be repeated weekly, up to 32 weeks' gestation.

Six- to 8-Year Follow-up

This mid-childhood follow-up of participating children was not part of the original protocol developed in 1997. Contact with families of surviving children was maintained both at the time of notification of the initial study results,² and the early childhood follow-up at 2 years of age.⁹ Parents gave written informed consent for their child to participate. The protocol for the school-age follow-up received human research ethics approval and is available in the supplemental material.

Surviving children were assessed at 6- to 8-years' corrected age by a pediatrician and a psychologist who were unaware of the child's randomized group. The pediatric assessment included a history

of health and health service use; physical and neurologic examination; measurement of weight, height, and head circumference; blood pressure; lung function; assessment of vision and hearing; and assessment of gross and fine motor function by using the Movement Assessment Battery for Children, Second Edition.¹⁰ Some children were assessed with an earlier edition of the Movement Assessment Battery for Children. Cerebral palsy included nonprogressive loss of motor function with disordered muscle tone or tendon reflexes.¹¹ The severity of gross motor function in children with cerebral palsy was assessed by using criteria of Palisano et al¹² and graded into mild (grade 1), moderate (grades 2-3), and severe (grades 4-5). Blindness comprised vision worse than 6/60 in the better eye. Deafness comprised hearing loss requiring hearing aids, or worse. Lung function was measured by portable flow spirometry (EasyOne 2001; NDD Technologies, Zurich, Switzerland) by using forced expiratory maneuvers. Spirogram quality was determined by 2 independent examiners, and children with at least 2 adequate and reproducible spirograms were included in the analysis.¹³

The psychological assessment included the Wechsler Abbreviated Scale of Intelligence.¹⁴ Full-scale IQ was derived from 4 subtests: Vocabulary, Similarities, Block Design, and Matrix Reasoning. Verbal and Performance IQ were also generated. Scores are age standardized (mean [M] = 100, SD = 15). Children unable to complete the Wechsler Abbreviated Scale of Intelligence because of severe intellectual impairment were assigned a score of 40. Intellectual impairment was classified as mild (IQ from -2 SD to <-1 SD), moderate (-3 SD to <-2 SD), or severe (<-3 SD).

Neurosensory disability comprised any of cerebral palsy, IQ <-1 SD, blindness or deafness, and was

graded as mild (mild cerebral palsy or mild intellectual impairment), moderate (deafness, moderate cerebral palsy, or moderate intellectual impairment), or severe (blindness, severe cerebral palsy, or severe intellectual impairment).

Academic skills were assessed by the word reading, spelling, and math computation subtests of the Wide Range Achievement Test.¹⁵ Each scale is age standardized ($M = 100$, $SD = 15$). Attention was assessed using subtests from the Test of Everyday Attention for Children.¹⁶

Selective visual attention was assessed with the Sky Search subtest, sustained attention with the Score! subtest, shifting attention with the Creature Counting subtest, and divided attention with the Sky Search Dual Task subtest. For the Sky Search, Score!, and Creature Counting subtests, performance was judged by age-standardized scores of accuracy ($M = 10$, $SD = 3$). For the Sky Search Dual Task, performance was determined by the average of (proportion of visual targets correctly identified plus proportion of correct auditory counting games) $\times 100$.¹⁶ Although there are no published norms for this scoring procedure, the range of possible values is 0 to 100, and in a study of 173 control children at 8 years of age the mean (SD) score was 80.3 (16.5).¹⁷

Attention-deficit/hyperactivity disorder (ADHD) symptoms were evaluated with parent and teacher versions of the Conners' ADHD/DSM-IV Scales (CADS; Psychological Corporation), which have age/sex t -scores ($M = 50$, $SD = 10$; higher scores indicate more problems).¹⁸

Executive function was assessed with the Rey Complex Figure,¹⁹ the Fruit Stroop Task,²⁰ and the Behavior Rating Inventory of Executive Function (BRIEF).²¹ The Rey Complex Figure assesses spatial organization. Children's copying of a complex geometrical figure was scored for accuracy (maximum score

36),²² and strategic organization.²³ The Fruit Stroop Task assesses impulse control, with performance judged by the number of correct responses in 45 seconds. The BRIEF is completed by parents and teachers and assesses behavioral manifestations of inattention and executive function. Performance was judged by the General Executive Composite, and the Metacognition and Behavioral Regulation Indices ($M = 50$, $SD = 10$; higher scores indicate more problems).

Memory and learning was assessed with the Rey Auditory Verbal Learning Test.²⁴ The Rey Auditory Verbal Learning Test comprises 5 trials in which the child recalls a single list of 15 words. Performance was assessed by the number of correct words recalled in trial 1, and over all 5 trials.

Visual perception was assessed by the Visual Discrimination, Figure Ground, and Visual closure subtests of the Test of Visual-Perceptual Skills.²⁵ Each subtest has age-standardized scores ($M = 10$, $SD = 3$).

General behavior was assessed using the Total Difficulties score from parent and teacher reports of the Strengths and Difficulties Questionnaire (SDQ), with possible scores ranging from 0 to 40: normal 0 to 13, borderline 14 to 16, and abnormal ≥ 17 .²⁶

Health-related quality of life was measured by the parent-completed Multiattribute Health Status classification system.²⁷ A Health Utility Index is obtained, ranging from 1 for perfect health to 0 for death. The parent-completed Australian Authorized Adaptation of the Child Health Questionnaire²⁸ provided assessments of the child's psychosocial health (score 0–600) and physical health (score 0–400), with higher scores indicating better health.

The primary outcomes were survival free of any neurosensory disability, and the categorization of neurosensory disability as none, mild, moderate, or severe. Secondary

outcomes were mortality; cerebral palsy; blindness or deafness; z -scores for height, weight, BMI, and head circumference; expiratory flows on lung function; blood pressure z -scores and proportions in the abnormal ranges; IQ; attention and executive function; memory and learning; visual perception; academic achievement; behavior; health service utilization and reason for use; general health; and health-related quality of life.

Statistical Analysis

Data were analyzed by using SAS software, version 9.3 (SAS Institute, Inc, Cary, NC). Normative data were used to calculate age- and sex-adjusted z -scores for body size²⁹ and blood pressure,³⁰ and sex- and height-adjusted z -scores for spirometry.³¹ Primary analyses compared primary and secondary outcomes between children with and without exposure to repeat betamethasone by using generalized linear models, adjusted for potential confounders (gestational age at trial entry, antepartum hemorrhage, and preterm prelabor rupture of membranes), and clustering of children from multiple pregnancy by using generalized estimating equations. Treatment effects are presented as risk ratio (RR) for binary and mean difference for continuous outcomes, with 95% confidence intervals (CIs). Two-tailed $\alpha < 0.05$ was considered statistically significant, with no adjustment for multiple comparisons. Estimates of study power are included in the study protocol (supplemental material).

RESULTS

Participant flow is outlined in Fig 1. Of the 1146 fetuses known to be alive at randomization, the mortality rate to mid-childhood was similar between groups; repeat corticosteroids (5.5%; 31/568) compared with the placebo group (6.1%; 35/578) (RR 0.90, 95% CI 0.55 to 1.46, $P = .67$, Fig 1). Of the




FIGURE 1
Flowchart of randomization, treatment, and mid-childhood follow-up of participants.

survivors, 21 had withdrawn from the study before this follow-up, leaving 528 in the repeat corticosteroids group, and 531 in the placebo group, about whom some data were available for 93% ($n = 493$) of the repeat corticosteroids group and 93% ($n = 495$) of the placebo group at mid-childhood. There were no substantial differences in the children eligible for follow-up between those with some data and those without any data (see the Supplemental Table 5). Of the 988 children followed, parental questionnaires were received for 98%, pediatric and psychological assessments for 89%, and teacher questionnaires for 71%, with assessment or response rates similar in the repeat corticosteroids and placebo groups.

The treatment groups were well balanced for important perinatal and sociodemographic variables (Table 1), with exceptions of higher rates of antepartum hemorrhage and lower

rates of preterm prelabor rupture of the membranes in the repeat corticosteroids group compared with the placebo group. Gestational ages at initial dose of corticosteroids, at trial entry, and at birth, and ages of the children when assessed were similar for both groups (Table 1).

Primary Outcomes

The rates of survival free of neurosensory disability at 6 to 8 years did not differ significantly in repeat corticosteroid and placebo groups (78% vs 77%, RR 1.00, 95% CI 0.94 to 1.08, $P = .89$) (Table 2). There were no significant differences between the groups in the distribution of the severity of the neurosensory disabilities, or in any of the individual components; intellectual impairment, blindness, deafness, or cerebral palsy (Table 2). There were 66 deaths after randomization, 31 in the repeat

corticosteroid group and 35 in the placebo group. The rates of combined outcomes of death or moderate disability and of death or severe disability were similar in both groups (Table 2).

Secondary Outcomes

Health Outcomes

There were no differences between the groups for body size, either as mean differences or proportions with measurements <10th percentile (Table 3).

The rates of asthma were equally high in both groups, as were the rates of requiring medication for asthma. Only 27% (185/691) of children tested could produce satisfactory expiratory flows. In those with valid data, there were no differences in expiratory flow rates between the groups (Table 3).

Mean systolic and diastolic blood pressure z-scores were not different between treatment groups, neither were the proportions of children with blood pressure in the prehypertensive range (Table 3). The rates of use of health services were also similar, with no differences in the need for readmission to hospital overall or for respiratory illnesses, or in the use of medical services (Table 3).

Functional Outcomes

There were no differences between groups in the rates of parent-assessed exercise tolerance (Table 3). Utilities and scores for the Child Health Questionnaire were similar between the treatment groups (Table 3).

Psychological Outcomes

There were no differences between the groups in Full-scale, Verbal, or Performance IQ (Table 4), or in scores for tests of attention and executive function, visual perception, or academic skills (Table 4). A small group difference (<0.2 SD) in favor of the placebo group was observed for the immediate recall of a list of 15 words, but no differences in recalling the words over 5 trials.

TABLE 1 Perinatal and Demographic Characteristics of Mothers and Children With Any Follow-up at Mid-Childhood

Variables	Repeat Steroids	Placebo
Mothers, <i>n</i>	430	426
Age at entry, y, mean (SD)	30.2 (6.0)	30.1 (5.8)
Gestation at first corticosteroids, median (25th–75th percentiles)	26.5 (24.5–28.5)	26.6 (24.6–28.5)
Gestation at entry, median (25th–75th percentiles)	28.4 (26.2–30.2)	28.4 (26.3–30.1)
Primiparous	32 (139)	32 (135)
Multiple pregnancy	16 (69)	16 (70)
Previous perinatal death	8 (35)	9 (37)
Previous preterm birth	13 (58)	16 (68)
Days from randomization to delivery, median (25th–75th percentiles)	25.5 (7.4–52.1)	23.0 (7.9–49.0)
Major reason for preterm birth		
Antepartum hemorrhage	33 (141)	26 (112)
Prelabor, preterm rupture of membranes	29 (125)	36 (152)
Preterm labor	27 (114)	24 (104)
Speak only English at home	87 (374)	87 (370)
Intact family	61 (263)	64 (274)
Socioeconomic status		
Home duties only	31 (132)	30 (129)
Lower	38 (162)	33 (142)
Higher	33 (142)	34 (144)
Doses of treatment given		
0	1 (4)	2 (7)
1	39 (169)	42 (178)
2–3	33 (143)	37 (158)
4	27 (114)	19 (83)
Infants, <i>n</i>	493	495
Gestational age at birth, completed wk, mean (SD)	32.5 (3.6)	32.4 (3.6)
Birth weight, g, mean (SD)	1917 (831)	1910 (808)
Boys	56 (275)	54 (268)
Mid-childhood follow-up		
Corrected age when assessed, y, mean (SD)	7.7 (1.1)	7.8 (1.2)

Data are % (*n*), unless otherwise specified.

TABLE 2 Primary Outcomes and Their Components at Mid-childhood

Outcomes	Repeat Steroids, <i>n</i> = 493	Placebo, <i>n</i> = 495	RR (95% CI) ^a	<i>P</i> ^a
Survival free of disability	78% (375/479)	77% (374/484)	1.00 (0.94 to 1.08)	.89
Death, <i>n</i>	31	35		
Neurosensory disability, <i>n</i>	448	449		.83 ^b
Nil	84 (375)	83 (374)		
Mild	12 (54)	12 (54)		
Moderate	2 (8)	2 (9)		
Severe	2 (11)	3 (12)		
IQ, mean (SD), <i>n</i>	99.9 (16.2), 444	99.7 (15.8), 445	0.0 (–2.2 to 2.3)	.99
IQ 70 to <85	11 (48)	11 (47)	1.04 (0.69 to 1.57)	.85
IQ 55 to <70	1 (5)	0.2 (1)	5.34 (0.64 to 44.5)	.12
IQ <55	2 (7)	2 (8)	0.89 (0.33 to 2.41)	.81
Blind	0.4 (2)	0.6 (3)	0.65 (0.11 to 3.91)	.64
Deaf	1.4 (7)	1.0 (5)	1.38 (0.43 to 4.41)	.59
Cerebral palsy	4 (19/493)	4 (20/494)	0.99 (0.54 to 1.82)	.97
Death or moderate/severe disability	10 (50/479)	12 (56/484)	0.92 (0.63 to 1.33)	.64
Death or severe disability	9 (42/479)	10 (47/484)	0.91 (0.61 to 1.39)	.67

Data are % (*n*), unless otherwise specified.

^a Adjusted for gestational age at entry, antepartum hemorrhage, prelabor, preterm rupture of membranes, and for clustering within mother.

^b From test for linear trend over all categories.

On the parent-rated perceptions of their child's behavior, there were no differences in scores on the BRIEF or Conners Scales, or in the SDQ raw scores (Table 4). Similarly, teacher rated perceptions of behavior did not differ between groups (Table 4).

DISCUSSION

In our study, repeat corticosteroids in women at risk for very preterm birth ≥ 7 days after an initial course reduced serious neonatal morbidity and respiratory disease² without

adverse effects on survival free of neurosensory disability, or on any other health or functional outcomes in mid-childhood. Given this evidence of longer-term safety, we consider that eligible women should receive repeat corticosteroids in view

TABLE 3 Health Outcomes at Mid-childhood.

Outcomes	Repeat Steroids, <i>n</i> = 493	Placebo, <i>n</i> = 495	Mean Difference/RR (95% CI) ^a	<i>P</i> ^a
Auxology				
Weight z-score	0.12 (1.27) 475	0.18 (1.36) 465	-0.06 (-0.24 to 0.12)	.52
Height z-score	0.15 (1.14) 461	0.13 (1.20) 451	0.02 (-0.15 to 0.18)	.85
BMI z-score	0.07 (1.27) 461	0.20 (1.29) 449	-0.14 (-0.32 to 0.04)	.14
Head circumference z-score	-0.49 (1.33) 442	-0.45 (1.36) 443	-0.04 (-0.24 to 0.14)	.62
Weight <10th percentile	12.2% (58/475)	13.0% (60/465)	0.96 (0.67 to 1.39)	.83
Height <10th percentile	10.6% (49/461)	11.3% (51/451)	0.94 (0.64 to 1.40)	.78
Head circumference <10th percentile	26.9% (118/442)	24.2% (107/443)	1.12 (0.89 to 1.41)	.34
Lung function				
FVC z-score	-0.04 (0.94) 98	0.02 (1.01) 87	-0.07 (-0.35 to 0.21)	.62
FEV ₁ z-score	-0.92 (1.10) 98	-0.74 (1.04) 87	-0.16 (-0.46 to 0.14)	.30
FEV ₁ /FVC z-score	-1.54 (0.85) 98	-1.35 (0.88) 87	-0.15 (-0.39 to 0.12)	.23
FEF _{25% to 75%} z-score	-1.55 (1.25) 98	-1.29 (1.19) 87	-0.21 (-0.55 to 0.12)	.21
Asthma ever	35.6% (174/489)	35.3% (173/490)	0.99 (0.84 to 1.18)	.94
Breathing medication ever	29.5% (144/488)	27.8% (136/489)	1.05 (0.85 to 1.29)	.66
Blood pressure				
Systolic - z-score	-0.26 (0.97) 423	-0.33 (0.96) 425	-0.09 (-0.26 to 0.08)	.32
Diastolic - z-score	-1.14 (1.21) 423	-1.05 (1.20) 425	0.07 (-0.07 to 0.20)	.34
Systolic or diastolic >90th percentile	7.1% (30/423)	6.4% (27/425)	1.07 (0.64 to 1.80)	.79
Use of health services				
Any hospital readmission	14% (67/489)	12% (61/291)	1.09 (0.78 to 1.53)	.62
Hospital readmission for respiratory illness	2% (11/489)	3% (14/291)	0.78 (0.36 to 1.72)	.56
Any visits to doctors	78% (377/485)	76% (374/491)	1.00 (0.93 to 1.08)	.91
Functional outcomes				
No limitation with physical activities requiring lots of energy	79% (373/470)	81% (378/465)	0.98 (0.91 to 1.04)	.50
Health Utility Index	0.904 (0.128) 470	0.910 (0.129) 467	-0.006 (-0.023 to 0.011)	.46
Child Health Questionnaire, Physical score	353.5 (59.0) 468	353.7 (60.1) 465	-0.5 (-8.7 to 7.6)	.90
Child Health Questionnaire, Psychosocial score	482.1 (95.7) 465	485.0 (90.5) 463	-2.6 (-15.3 to 10.1)	.68

Data are % (*n*), number with data, or mean (SD), number with data, unless otherwise specified. From test for linear trend over all categories. FEF, forced expiratory flow; FEV₁, forced expiratory volume in 1 second; FVC, forced vital capacity.

^a Adjusted for gestational age at entry, antepartum hemorrhage, prelabor, preterm rupture of membranes, and for clustering within mother.

of the clinically important neonatal benefits.

Despite widespread use of beneficial, single-course antenatal glucocorticoids,³² very preterm infants continue to have high neonatal morbidity. In our trial, control infants, all of whom were exposed to a single course of antenatal corticosteroids, frequently developed respiratory distress syndrome (RDS) (41%), severe lung disease (20%), and other serious complications related to preterm birth (26%).² Treatment with repeat dose(s) of antenatal corticosteroid substantially reduced these complications (RDS RR [95% CI] 0.82 [0.71 to 0.95]; severe lung disease 0.60 [0.46 to 0.79]; serious neonatal morbidity 0.79 [0.65 to 0.97]). Further, the absolute benefit was similar to that achieved with a single course (number needed to treat [95% CI] to prevent 1 case of RDS: 14 [8 to 50] for repeat dose(s)²; 12 [7 to 14] for a single course³²).

This suggests that serial antenatal corticosteroid exposure is required for maximal maturational response in the preterm fetus. By reducing the severity of neonatal morbidities, it is likely that repeat antenatal corticosteroid treatment also reduces health care costs.

Despite evidence of early benefit, repeat antenatal corticosteroids are not commonly prescribed, largely because of concern about possible longer-term adverse effects, particularly on neurologic development.^{33,34} In early childhood in our trial, children exposed to repeat corticosteroids were more likely to have scores in the clinical range for attention problems at 2 years' corrected age compared with those exposed to a single course, although no other early childhood outcomes differed between groups.⁹ Our results in mid-childhood show that early exposure to repeat corticosteroids

does not have any adverse effect on later neurocognitive function, including attention and behavior, with 1 minor exception, suggesting that our earlier finding of an effect on attention scores may have been a type 1 error or reflected a specific developmental delay. Our data are robust, with similar scores and rates of impairment in both groups using an extensive array of assessments, including psychometric tests across multiple domains, including general cognitive ability, executive function, attention, memory and learning, visual-perceptual skills, behavior, and educational achievement. There were also no differences between groups in motor performance and rates of neuromotor impairment. By mid-childhood, cognitive assessments are highly predictive of later function,³⁵ unlike preschool assessment of cognitive functioning.³⁶ Therefore, we consider there is now strong evidence that repeat doses of antenatal

TABLE 4 Cognitive, Academic, Attention, Executive Function, and Behavioral Outcomes at Mid-childhood.

Outcome	Repeat Steroids	Placebo	Mean Difference/RR (95% CI) ^{a, b}	P ^{a, b}
Cognitive				
Full-scale IQ	99.9 (16.2) 444	99.7 (15.8) 445	0.0 (−2.2 to 2.3)	.99
Verbal IQ	100.3 (15.5) 440	101.2 (15.1) 438	−1.3 (−3.5 to 0.9)	.26
Performance IQ	101.5 (14.8) 438	100.4 (13.9) 437	1.0 (−1.0 to 3.1)	.34
Academic skills (WRAT-4)				
Reading	99.1 (16.6) 423	99.2 (17.9) 427	−0.2 (−2.7 to 2.4)	.91
Spelling	99.7 (15.6) 420	100.5 (16.8) 426	−0.9 (−3.4 to 1.6)	.48
Mathematics	96.8 (15.8) 419	96.5 (16.0) 426	0.1 (−2.3 to 2.4)	.97
Attention				
Selective – Sky Search	9.2 (3.1) 426	9.1 (3.3) 428	0.03 (−0.4 to 0.5)	.89
Sustained – Score!	8.7 (3.5) 422	8.8 (3.6) 408	−0.1 (−0.6 to 0.5)	.84
Shifting – Creature Counting	9.7 (3.7) 377	9.5 (3.6) 370	0.2 (−0.4 to 0.7)	.52
Divided – Sky Search Dual Task	58.9 (29) 408	59.8 (29.4) 390	−0.8 (−5.0 to 3.4)	.71
Executive function				
Rey Complex Figure				
Copy score	15.5 (7.7) 426	15.3 (7.9) 422	0.4 (−0.5 to 1.4)	.39
Organization	3.7 (1.2) 424	3.6 (1.2) 421	0.1 (−0.04 to 0.3)	.16
Fruit Stroop – Trial 4, number correct	19.5 (8.3) 413	20.8 (8.4) 416	−0.9 (−2.0 to 0.1)	.08
BRIEF – Parent				
Global Executive Composite	52.5 (13.1) 428	52.1 (12.5) 429	0.3 (−1.6 to 2.1)	.79
Metacognition Index	52.3 (13.0) 428	52.0 (12.7) 430	0.3 (−1.6 to 2.1)	.78
Behavioral Regulation Index	51.9 (12.9) 428	51.5 (12.2) 429	0.3 (−1.6 to 2.1)	.77
BRIEF – Teacher				
Global Executive Composite	53.6 (12.4) 349	55.4 (13.5) 342	−1.9 (−4.1 to 0.3)	.08
Metacognition Index	54.7 (13.4) 349	56.2 (13.9) 341	−1.6 (−3.8 to 0.6)	.14
Behavioral Regulation Index	51.7 (11.5) 351	53.6 (13.5) 343	−2.0 (−4.1 to 0.1)	.05
Memory and Learning - Rey Auditory Verbal Learning Test				
Words correct trial 1	4.5 (1.8) 428	4.8 (1.9) 429	−0.3 (−0.5 to −0.1)	.01
Words correct all trials	37.1 (11.1) 427	37.5 (11.3) 428	−0.3 (−1.8 to 1.2)	.71
Visual-Perceptual Skills				
Visual discrimination	9.0 (3.5) 424	8.6 (3.6) 428	0.3 (−0.2 to 0.8)	.24
Figure ground	10.3 (4.1) 421	10.2 (4.2) 429	0.1 (−0.5 to 0.7)	.72
Visual closure	8.7 (3.9) 421	8.5 (3.8) 428	0.3 (−0.3 to 0.8)	.36
Behavior				
CADS – Parent t-scores				
ADHD Index	51.1 (6.7) 432	51.5 (7.0) 435	−0.4 (−1.3 to 0.6)	.43
DSM-IV Inattentive	51.2 (7.6) 432	51.4 (7.1) 435	−0.2 (−1.3 to 0.8)	.65
DSM-IV Hyperactive-impulsive	52.0 (7.2) 432	52.6 (7.0) 435	−0.8 (−1.8 to 0.2)	.13
DSM-IV Total	51.6 (7.2) 432	52.1 (6.9) 435	−0.5 (−1.5 to 0.5)	.29
CADS – Teacher t-scores				
ADHD Index	49.4 (6.6) 351	50.5 (7.4) 345	−1.1 (−2.2 to −0.0)	.06
DSM-IV Inattentive	46.8 (5.7) 351	47.1 (5.7) 345	−0.4 (−1.3 to 0.5)	.35
DSM-IV Hyperactive-impulsive	49.4 (6.8) 351	50.1 (7.0) 345	−0.8 (−1.9 to 0.3)	.15
DSM-IV Total	49.2 (6.3) 351	49.8 (6.3) 345	−0.6 (−1.6 to 0.3)	.24
SDQ Total Difficulties				
– Parent	11.0 (6.9) 431	10.7 (6.4) 434	0.3 (−0.7 to 1.3)	.53
– Teacher	9.2 (6.7) 353	9.8 (7.2) 345	−0.7 (−1.8 to 0.4)	.22

Data are mean (SD), number with data, unless otherwise specified. DSM-IV, *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition*; WRAT, Wide Range Achievement Test.

corticosteroids do not adversely affect long-term neurologic development.

Investigators from the Multiple Courses of Antenatal Corticosteroids Study (MACS) recently reported behavior and developmental outcomes for survivors at 5 years of age by using a parental questionnaire, with formal cognitive assessment completed on <20% of children.³⁷ They also found no

differences in rates of death or severe disability between groups (repeat corticosteroids 25% [217/817]; controls 25% [201/848]; odds ratio 1.02, 95% CI 0.81 to 1.29; $P = .84$), and no differences in body size or blood pressure. They did not report on lung function, and we recognize that the inability to obtain adequate lung function data on most children assessed is a limitation of our study.

Another important concern has been whether exposure to repeat doses of corticosteroids could result in long-term perturbations of metabolic and cardiovascular function, increasing the risk of adult disease, as seen in several animal studies.^{6,7} In our trial, repeat corticosteroid treatment was associated with a temporary slowing of fetal growth,² a potential marker of fetal programming,³⁸ although the

effect was small and short-lived,² being of similar magnitude to that seen in infants born ≥ 1 days after a single course of corticosteroids.³² However, we found no evidence of altered blood pressure, body size, or general health at mid-childhood in those exposed to repeat compared with single courses of antenatal corticosteroids. Further, our in-depth side-study has reported no increase in physiologic risk factors for cardiovascular or metabolic disease at early school age.³⁹

Although our trial sample was representative of women at ongoing risk of preterm birth after an initial course of corticosteroids,⁴⁰ caution is required in generalizing our results to other treatment protocols or inclusion criteria. The recent Cochrane systematic review included 9 other randomized trials of repeat antenatal corticosteroid treatment, with varying number, frequency, and dose(s) of corticosteroids.¹ Although the meta-analysis confirmed that repeat antenatal corticosteroid treatment is associated with neonatal benefits,¹ broadly similar to those seen in our trial, some heterogeneity of effect was observed, including 2 trials that showed no apparent benefit.^{4,5} Until this heterogeneity is adequately explained and later childhood outcomes are reported for different treatment regimens, we recommend that the inclusion criteria and treatment schedule used in our trial be followed. An important feature of our protocol was that the risk of preterm birth was reevaluated on a weekly basis, so that most women received only 1 or 2 repeat doses.

CONCLUSIONS

We have shown that the use of repeat dose(s) of antenatal betamethasone when indicated reduces neonatal morbidity without adverse effects on survival free of neurosensory disability in mid-childhood, and no other apparent harmful or beneficial effects. For women at risk for preterm birth before 32 weeks'

gestation, ≥ 7 days after an initial course of antenatal corticosteroids, clinicians could consider using a single injection of betamethasone, repeated weekly if risk remains.

ACKNOWLEDGMENTS

We thank all the women and their children who participated in this study.

Members of the ACTORDS Follow-up Group who collaborated to acquire data for the 6- to 8-year follow-up study are listed as follows.

Central coordinating team University of Adelaide: Caroline Crowther, Pat Ashwood, Vincent Ball, Amy Earl, Carol Holst, Kaye Robinson, Yu Zhang.

Statistical support: Dr Kristyn Willson was the trial biostatistician and conducted the statistical analyses.

The collaborating centers where children were assessed and the participating investigators were as follows (the total number of children assessed at each center is given in parentheses).

Canberra Hospital, Australian Capital Territory, Australia (28): Graham Reynolds, Pat Ashwood, Rachel Earl, Jan Holmes, Manina Pathak, Sophie Trenowden, Sue Wrigley.

Liggins Institute, University of Auckland, New Zealand (307): Jane Harding, Chris McKinlay, Coila Bevan, Janet Holmes, Judith Ansell, Malcolm Battin, Stuart Dalziel.

Mater Mothers' Hospital, Brisbane, Queensland, Australia (60): Peter Gray, Vicki Flenady, Peter Gray, David Hou, Elizabeth Hurrion, Luke Jardine, Lisa Lynn, Leith Poulsen, David Tudehope.

Mercy Hospital for Women, Victoria, Australia (60): Dan Casalaz, Noni Davis, Elaine Kelly, Justin Kelly, Emma Magrath, Emma Marchant, Gillian Opie, Anne-Marie Turner, Amanda Williamson, Heather Woods.

Monash Medical Centre, Victoria, Australia (24): Elizabeth Carse, Margaret Charlton, Marie Hayes, Elaine Kelly, Kerry Saunders.

Royal Brisbane and Women's Hospital, Queensland, Australia (62): Paul Colditz, David Cartwright, Brooke Dougan, Anna Pritchard, Margo Pritchard.

Royal Hobart Hospital, Tasmania, Australia (11): Antonio De Paoli, Pat Ashwood, Rachel Earl.

Royal North Shore Hospital, New South Wales, Australia (41): Jennifer Bowen, Linda Hayes-Cameron, Mary Paradisis, Himanshu Popat, Sara Sedgley.

Royal Women's Hospital, Victoria, Australia (78): Lex Doyle, Catherine Callanan, Noni Davis, Cinzia Deluca, Julianne Duff, Esther Hutchinson, Elaine Kelly, Louise Kornman, Marion McDonald, Gehan Roberts.

Townsville Hospital, Queensland, Australia (24): Koh Thhg, Pat Ashwood, Rachel Earl, Kim Pfitzner, Pat Ryan, Sophie Trenowden.

Women's and Children's Hospital, South Australia, Australia (201): Caroline Crowther, Chad Andersen, Pat Ashwood, Vineesh Bhatia, Amy Earl, Rachel Earl, Louise Goodchild, Freya Goodhew, Karen Harris, Ross Haslam, Bevan Headley, Rosslyn Lontis, Andrew McPhee, Kim Pfitzner, Jeffrey Robinson, Michael Stark, Laura Tisato, Sophie Trenowden.

ABBREVIATIONS

ACTORDS: The Australasian Collaborative Trial of Repeat Doses of Corticosteroids

ADHD: attention-deficit hyperactivity disorder

BRIEF: Behavior Rating Inventory of Executive Function

CADS: Connors ADHD/DSM-IV Scales

CI: confidence interval

M: mean

RDS: respiratory distress syndrome

RR: risk ratio

SDQ: Strengths and Difficulties Questionnaire

participated in data interpretation; Dr Doyle conceptualized and designed the study, and participated in data acquisition, analyses, and interpretation; and all authors revised the manuscript and approved the final manuscript as submitted.

This trial has been registered with the ACTRN Register (Identifier ACTRN12606000318583).

DOI: 10.1542/peds.2016-0947

Accepted for publication Jul 15, 2016

Address correspondence to Caroline Crowther, MD, Liggins Institute, The University of Auckland, Private Bag 92019, 85 Park Rd, Grafton, Auckland, New Zealand. E-mail: c.crowther@auckland.ac.nz

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2016 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: This follow-up study was supported through a 4-year project grant from the Australian National Health and Medical Research Council (453633), a 3-year project grant from the New Zealand Health Research Council (07/204), and a 2-year project grant from the Auckland Medical Research Foundation (81537).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

1. Crowther CA, McKinlay CJD, Middleton P, Harding JE. Repeat doses of prenatal corticosteroids for women at risk of preterm birth for improving neonatal health outcomes. *Cochrane Database Syst Rev.* 2015;(7):CD003935
2. Crowther CA, Haslam RR, Hiller JE, Doyle LW, Robinson JS; Australasian Collaborative Trial of Repeat Doses of Steroids (ACTORDS) Study Group. Neonatal respiratory distress syndrome after repeat exposure to antenatal corticosteroids: a randomised controlled trial. *Lancet.* 2006;367(9526):1913–1919
3. Wapner RJ, Sorokin Y, Thom EA, et al; National Institute of Child Health and Human Development Maternal Fetal Medicine Units Network. Single versus weekly courses of antenatal corticosteroids: evaluation of safety and efficacy. *Am J Obstet Gynecol.* 2006;195(3):633–642
4. Peltoniemi OM, Kari MA, Tammela O, et al; Repeat Antenatal Betamethasone Study Group. Randomized trial of a single repeat dose of prenatal betamethasone treatment in imminent preterm birth. *Pediatrics.* 2007;119(2):290–298
5. Murphy KE, Hannah ME, Willan AR, et al; MACS Collaborative Group. Multiple courses of antenatal corticosteroids for preterm birth (MACS): a randomised controlled trial. *Lancet.* 2008;372(9656):2143–2151
6. de Vries A, Holmes MC, Heijnis A, et al. Prenatal dexamethasone exposure induces changes in nonhuman primate offspring cardiometabolic and hypothalamic-pituitary-adrenal axis function. *J Clin Invest.* 2007;117(4):1058–1067
7. Moss TJM, Doherty DA, Nitsos I, Sloboda DM, Harding R, Newnham JP. Effects into adulthood of single or repeated antenatal corticosteroids in sheep. *Am J Obstet Gynecol.* 2005;192(1):146–152
8. French NP, Hagan R, Evans SF, Mullan A, Newnham JP. Repeated antenatal corticosteroids: effects on cerebral palsy and childhood behavior. *Am J Obstet Gynecol.* 2004;190(3):588–595
9. Crowther CA, Doyle LW, Haslam RR, Hiller JE, Harding JE, Robinson JS; ACTORDS Study Group. Outcomes at 2 years of age after repeat doses of antenatal corticosteroids. *N Engl J Med.* 2007;357(12):1179–1189
10. Henderson SE, Sugden DA, Barnett A. *Movement Assessment Battery for Children.* 2nd ed. Sidcup, Kent: Harcourt Assessment; 2007
11. Kitchen WH, Doyle LW, Ford GW, Rickards AL, Lissenden JV, Ryan MM. Cerebral palsy in very low birthweight infants surviving to 2 years with modern perinatal intensive care. *Am J Perinatol.* 1987;4(1):29–35
12. Palisano R, Rosenbaum P, Walter S, Russell D, Wood E, Galuppi B. Development and reliability of a system to classify gross motor function in children with cerebral palsy. *Dev Med Child Neurol.* 1997;39(4):214–223
13. American Thoracic Society. Standardization of spirometry, 1994 update. *Am J Respir Crit Care Med.* 1995;152(3):1107–1136
14. Wechsler D. *The Wechsler Abbreviated Scale of Intelligence.* San Antonio, TX: The Psychological Corporation; 1999
15. Wilkinson G, Robertson G. *Wide Range Achievement Test,* 4th ed. Wilmington, DE: Pearson; 2005
16. Manly T, Robertson I, Anderson VA, Nimmo-Smith I. *The Test of Everyday Attention for Children.* Suffolk, UK: Thames Valley Test Company; 1999
17. Anderson PJ, De Luca CR, Hutchinson E, Spencer-Smith MM, Roberts G, Doyle LW; Victorian Infant Collaborative Study Group. Attention problems in a representative sample of extremely preterm/extremely low birth weight children. *Dev Neuropsychol.* 2011;36(1):57–73
18. Conners CK. *Conners ADHD/DSM-IV Scales (CADS).* North Tonawanda, NY: Multi-Health Systems; 1997
19. Rey A. Psychological examination of traumatic encephalopathy (translation). *Clin Neuropsychol.* 1993;7(1):3–21
20. Archibald SJ, Kerns KA. Identification and description of new tests of executive functioning in children. *Child Neuropsychol.* 1999;5(2):115–129
21. Gioia G, Isquith P, Guy S, Kenworthy L. *BRIEF: Behavior Rating Inventory of Executive Function. Professional Manual.* Odessa, FL: Psychological Assessment Resources; 2000

22. Osterrieth P. The complex figure copy test (translation). *Clin Neuropsychol*. 1993;7(1):3–21
23. Anderson P, Anderson V, Garth J. Assessment and development of organizational ability: the Rey Complex Figure Organizational Strategy Score (RCF-OSS). *Clin Neuropsychol*. 2001;15(1):81–94
24. Rey A. *L'Examen Clinique en Psychologie*. Paris, France: Presses Universitaires de France; 1964
25. Martin NA. *Test of Visual-Perceptual Skills (No-Motor)*, 3rd ed. Novato, CA: Academic Therapy Publications; 2006
26. Goodman R. The strengths and difficulties questionnaire: a research note. *J Child Psychol Psychiatry*. 1997;38(5):581–586
27. Saigal S, Rosenbaum P, Stoskopf B, et al. Comprehensive assessment of the health status of extremely low birth weight children at eight years of age: comparison with a reference group. *J Pediatr*. 1994;125(3):411–417
28. Waters E, Salmon L, Wake M, Hesketh K, Wright M. The Child Health Questionnaire in Australia: reliability, validity and population means. *Aust N Z J Public Health*. 2000;24(2):207–210
29. Cole TJ, Williams AF, Wright CM; RCPCH Growth Chart Expert Group. Revised birth centiles for weight, length and head circumference in the UK-WHO growth charts [published correction appears in *Ann Hum Biol*. 2011;38(1):7–11]. *Ann Hum Biol*. 2011;38(2):241. *Ann Hum Biol*. 2011;38(1):7–11
30. Jackson LV, Thalange NKS, Cole TJ. Blood pressure centiles for Great Britain. *Arch Dis Child*. 2007;92(4):298–303
31. Stanojevic S, Wade A, Stocks J, et al. Reference ranges for spirometry across all ages: a new approach. *Am J Respir Crit Care Med*. 2008;177(3):253–260
32. Roberts D, Dalziel S. Antenatal corticosteroids for accelerating fetal lung maturation for women at risk of preterm birth. *Cochrane Database Syst Rev*. 2006;(3):CD004454
33. Quinlivan JA, Beazley LD, Braekevelt CR, Evans SF, Newnham JP, Dunlop SA. Repeated ultrasound guided fetal injections of corticosteroid alter nervous system maturation in the ovine fetus. *J Perinat Med*. 2001;29(2):112–127
34. Modi N, Lewis H, Al-Naqeeb N, Ajayi-Obe M, Doré CJ, Rutherford M. The effects of repeated antenatal glucocorticoid therapy on the developing brain. *Pediatr Res*. 2001;50(5):581–585
35. Breslau N, Johnson EO, Lucia VC. Academic achievement of low birthweight children at age 11: the role of cognitive abilities at school entry. *J Abnorm Child Psychol*. 2001;29(4):273–279
36. Roberts G, Anderson PJ, Doyle LW; Victorian Infant Collaborative Study Group. The stability of the diagnosis of developmental disability between ages 2 and 8 in a geographic cohort of very preterm children born in 1997. *Arch Dis Child*. 2010;95(10):786–790
37. Asztalos EV, Murphy KE, Willan AR, et al; MACS-5 Collaborative Group. Multiple courses of antenatal corticosteroids for preterm birth study: outcomes in children at 5 years of age (MACS-5). *JAMA Pediatr*. 2013;167(12):1102–1110
38. Barker DJP. The developmental origins of well-being. *Philos Trans R Soc Lond B Biol Sci*. 2004;359(1449):1359–1366
39. McKinlay CJD, Cutfield WS, Battin MR, Dalziel SR, Crowther CA, Harding JE; ACTORDS Study Group. Cardiovascular risk factors in children after repeat doses of antenatal glucocorticoids: an RCT. *Pediatrics*. 2015;135(2). Available at: www.pediatrics.org/cgi/content/full/135/2/e405
40. McLaughlin KJ, Crowther CA, Vigneswaran P, Hancock E, Willson K. Who remains undelivered more than seven days after a single course of prenatal corticosteroids and gives birth at less than 34 weeks? *Aust N Z J Obstet Gynaecol*. 2002;42(4):353–357

Mid-Childhood Outcomes of Repeat Antenatal Corticosteroids: A Randomized Controlled Trial

Caroline A. Crowther, Peter J. Anderson, Christopher J.D. McKinlay, Jane E. Harding, Pat J. Ashwood, Ross R. Haslam, Jeffery S. Robinson, Lex W. Doyle and for the ACTORDS Follow-up Group
Pediatrics 2016;138;; originally published online September 20, 2016;
DOI: 10.1542/peds.2016-0947

Updated Information & Services	including high resolution figures, can be found at: /content/138/4/e20160947.full.html
Supplementary Material	Supplementary material can be found at: /content/suppl/2016/09/17/peds.2016-0947.DCSupplemental.html
References	This article cites 31 articles, 5 of which can be accessed free at: /content/138/4/e20160947.full.html#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Fetus/Newborn Infant /cgi/collection/fetus:newborn_infant_sub Gynecology /cgi/collection/gynecology_sub Maternal and Fetal Medicine /cgi/collection/maternal_fetal_medicine_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: /site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: /site/misc/reprints.xhtml

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Mid-Childhood Outcomes of Repeat Antenatal Corticosteroids: A Randomized Controlled Trial

Caroline A. Crowther, Peter J. Anderson, Christopher J.D. McKinlay, Jane E. Harding, Pat J. Ashwood, Ross R. Haslam, Jeffery S. Robinson, Lex W. Doyle and for the ACTORDS Follow-up Group
Pediatrics 2016;138;; originally published online September 20, 2016;
DOI: 10.1542/peds.2016-0947

The online version of this article, along with updated information and services, is located on the World Wide Web at:
</content/138/4/e20160947.full.html>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

